

Protocol 1000 Plus

This protocol is simply adding DMSO to the MMS1 Protocol 1000 dose which is 3 activated drops per hour (or sometimes less). With Protocol 1000 Plus, you add in 3 drops of DMSO for each drop of MMS. Or, in case you have not progressed to 3 drops an hour by this time, continue the same amount of MMS1 you've been taking and add in the DMSO accordingly. For example, if you are taking a 3-drop dose of MMS1, you would add 9 drops of DMSO. If you are taking a 2-drop dose of MMS1, you would add 6 drops of DMSO.

You can start using DMSO if you cannot go further than your current dose of 1, 2 or 3 drops, but still feel you need to improve.

Step 1

Activate 3 drops of MMS. (If you have not been able to work up to a 3-drop dose yet, due to nausea, etc. activate however many drops you are taking).

Add 120 ml of water.

Step 2

Immediately after adding the water, add in 3 drops of DMSO for each drop of MMS1 you are using. For example, if you are making a 3-drop MMS1 dose, add 9 drops of DMSO. (Thoroughly mix in the DMSO by stirring it with a plastic spoon).

Drink down the dose immediately as once the DMSO is added the MMS1 will begin to slowly lose potency if left to sit.

Step 3

If after adding DMSO to your dose you experience discomfort (such as nausea, diarrhea, etc.), reduce the amount of DMSO you are adding on the next dose. Instead of adding 3 drops of DMSO per 1 drop of MMS1, reduce to 2 drops of DMSO to 1 drop of MMS1.

If you still have discomfort, reduce the DMSO by another drop, in other words, use 1 drop of DMSO per 1 drop of MMS1.

If you still experience discomfort after reducing the amount of DMSO 2 times, then completely stop adding DMSO to your MMS1 doses for a day. Then start back with small doses of DMSO and build up slowly to 3 drops of DMSO per 1 drop of MMS1.

Protocol 3000 - Topical Application

The goal with serious or life-threatening situations is to quickly get MMS1 circulating in the blood while trying to stay under the nausea level. One way to achieve this is by using DMSO with MMS1 topically.

Step 1

Mix up a solution of 10 drops of MMS1 with 10 drops of 10 drops of Avtivor. Count 30 seconds for activation.

Add 20 drops of water. But if this mixture is too strong (causes burning of the skin or other irritation), add more drops of water until it doesn't cause irritation.

If there is no extra skin sensitivity, you may want to add less water (do this in increments), to see if you can tolerate a stronger mixture.

Add 1 teaspoon/5 ml of DMSO.

Immediately spread the mixture over one arm. You can use your hand to spread the mix. (Be sure your hands are washed and thoroughly rinsed so as to remove all of the soap before applying DMSO.) It is not necessary, and even potentially dangerous to wear a glove, (if latex or rubber). When you have finished, wash your hand with plain water, do not wash with soap and water, as DMSO is a carrier and can carry some of the soap into your tissues.

Step 2

The following hour, mix up another MMS1/DMSO solution and spread it on your other arm. Repeat the next hour and do a different part of your body. Do one arm first, then the other arm, then a leg, then the other leg, then your stomach, and then back to the first arm, and so on. Use a different part of the body each time you apply the DMSO/MMS1 combination. Do this once every hour for eight consecutive hours.

Step 3

Repeat this process once every hour for eight consecutive hours for three consecutive days. (It is fine to bathe or shower after completing the hours applying the MMS1/DMSO. Wait at least one hour after applying the last application. Thoroughly rinse well with plain water before using soap).

Step 3

Then take a break. Quit from one to four days or however many days it takes to overcome any problems that may be caused by the DMSO (such as extra dry skin). After the first week you can use this topical application four days a week, or more, if there is no problem with your skin. If there is no problem, continue to use MMS1 and DMSO every day, as long as there is no skin irritation. Anytime you experience irritation, cut back for a time, or you can add more water to the mixture.